

Short Communication

Open Access

Simplified Method for the Cultivation of Spirulina for Domestic Use

P. Srivani¹, K. Tony¹, S. Akhila¹, M. Nageswara Rao¹, M. Amaze¹, B. Rajesh¹, K. Vimila¹, K. Dhanalakshmi¹, K. Pavan Kalyan¹, M. Indira¹, Md. N. Bobby¹, S. Krupanidhi^{*1}, K. Prakash Narayana Reddy¹, R. Sivakumar²

¹Department of Biotechnology, Vignan's Foundation for Science, Technology and Research University, Vadlamudi 522213, Andhra Pradesh, India

²Tejas Biotech Pvt. Ltd, Chennai, India

Corresponding Author: S. Krupanidhi, Department of Biotechnology, Vignan's Foundation for Science, Technology and Research University, Vadlamudi 522213, Andhra Pradesh, India. Tel: +91 7980323577, E-mail: krupanidhi.srirama@gmail.com

Citation: S. Krupanidhi et al. (2017), Simplified Method for the Cultivation of Spirulina for Domestic Use. Int J Biotech & Bioeng. 3:4, 107-108. DOI: [10.25141/2475-3432-2017-4.0107](https://doi.org/10.25141/2475-3432-2017-4.0107)

Copyright: ©2017 S. Krupanidhi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Received: May 03, 2017; **Accepted:** May 17, 2017; **Published:** May 29, 2017

Abstract:

Spirulina is a blue-green algae. There are three predominant species in India namely *Spirulina platensis*, *Spirulina fusiformis* and *Spirulina maxima*. All of them are profusely enriched with nutrients. As a consequence, *Spirulina* formulations are made available in the market for use. Unique attributes of *Spirulina* are autotrophic and alkaliphyletic and hence the energy requirements to maintain its culture at home are almost nullified. The sunlight that we obtain at our homes is more than adequate for its growth and replication. Mineral replenishment only needs to be supplemented. Yet, another important ingredient in the culture medium is pH. *Spirulina* prefers to grow in extreme alkaline medium under sunlight that restrains from most of the facultative bacteria and fungi which otherwise contaminate the culture medium. Therefore, little sophistication is required for its maintenance at domestic premises. The medium composition tried in our lab led to the profuse growth of the culture as indicated through the intensity of the green colour in the wider plastic tubs and obtained biomass was processed for the preparation of dried flakes

Keywords: Spirulina Maxima, Mineral Composition, Cultivation, Spirulina Culture, In-House

Among several algal species, *Spirulina* was heavily exploited and became a commercially important filamentous cyanobacterium due to its inherent ability to accumulate wealth of natural resources such as proteins, minerals, PUFAs and a few vitamins. Hence, several methods are attempted to cultivate both at laboratory, mass and industrial scales (1, 2, 3). Outdoor pond algal cultivation for the production of single cell proteins is one of the oldest technologies intuitively initiated by Borowitzka and Borowitzka (2). Various attempts such as closed system (4), outdoor open tubs with paddle wheel (5), and photo-reactor system (6), designs using confectionary, agricultural and poultry waste (1) are in vogue. In one instance, the yield containing 35 tonnes of *Spirulina* per hectare per year from a commercial pond was achieved in the Siam Algae Site near Bangkok (7). The attributes of its mass cultivation primarily are due to its photosynthetic pigment, phycocyanin, adaptability, plasticity and its asexual reproduction by binary fission. Due to its increasing popularity as an abundant nutritional biological resource, it is being used in poultry, aquaculture, animal feed, wastewater treatment and agriculture (7 & 8) and also supplementing human food requirements. As a result the single cell protein of *Spirulina* is being recycled in the ecosystem at different trophic levels. Therefore keeping in view of its importance, it is

aimed at devising a simplest protocol to culture *Spirulina* at our residences being tropical climatic conditions, conducive for algal growth.

The mother culture of *Spirulina maxima* was obtained from M/S Tejas Biotech Pvt Ltd, Chennai. 50 ml mother culture was mixed initially with 200 ml designed culture medium. The designed medium composition was: NaCl:2.0 g., NaHCO₃:16.0 g., NaNO₃:2.0 g., K₂SO₄: 1.0 g., KH₂PO₄:0.5 g., FeSO₄: 100 mg per litre water (boiled and chilled) with pH value of 10.5. The initial culture was kept in an orbital shaker with natural illumination for 7 days and observed the intensity of green colour through UV spectrophotometer at 540 nm. Once, the OD value 0.8 was reached, the initial culture was transformed to 10.0 L volume plastic tubs for mass culture in 1:4 proportions (initial grown culture: medium). The tubs with mass culture were allowed under direct sunlight for 7 days. The growth of *Spirulina* was noticed everyday by observing the intensity of green colour and replenished with 2 L fresh medium on 4th day. At the end of the 7th day, harvest was done through a muslin cloth (Fig.1.8), slurry mass was centrifuged and then taken into a syringe barrel. With the help of the plunger, the semisolid *Spirulina* mass pushed through a narrow nozzle of the syringe and allowed to collect on a clean

porcelain tiled surface, de-moisturized under ceiling fan initially and later dried under sunlight. Dried Spirulina fragments appeared like green fragile needles which were ready for consumption. A few more observations were made such as growth of Spirulina culture and its microscopic morphology.



Figure 1: (1) Initial culture with 50 ml mother culture +150 ml medium, (2) 5th day of initial culture showing more green intensity, (3) 7th day initial culture mixed in 1: 4 proportion in 1.0 L medium in a open tub, (4) 0-day mass culture under direct sunlight, (5) 3rd day of mass culture with relatively good growth, (6) 7th day of mass culture, (7) Spirulina 10x magnification isolated from a dense mass culture, (8) Harvested cultured Spirulina using muslin cloth, (9) Harvested biomass centrifuged and (10) Obtained biomass was squeezed through a syringe as needles for drying under sunlight.

With the intention of developing a simple protocol for the housewife to adopt the technique of Spirulina culture at home to practice, we followed the aforementioned procedure described. There was a 4 fold increase in the biomass within 14 days of culture duration. To begin with 50 ml mother culture obtained from M/S Tejas Biotech Ltd, Chennai containing 0.738 mg was seeded into 200 ml culture medium (Fig.1.1 & 2) At the end of a series of

steps described under material and methods and as shown in Fig.1, there was a profuse growth (Fig.6) of pure culture of Spirulina (Fig.7) and we harvested 2.871 g (Fig.1.9). This slurry biomass was washed twice to bring pH to 7.2. The same was allowed to dry under direct sunlight for three hours after spreading as thin rods on a porcelain tile using a plastic syringe (Fig.1.10). The dried Spirulina was kept in an eppendorf vial for consumption. All the steps followed in the protocol did not require any laboratory sophistication. Further, this protocol yielded pure strains without any contamination (Fig.1.7). In addition, the inorganic chemicals used are neither rare nor unaffordable. Industrial scale cultures are being practiced which requires optimization both at the level of the maintenance of medium concentration and also prevention of facultative pathogens (3).

References

- 1.Hala Y. El-Kassas , Ahmed M.M. Heneash and Nabila R. Hussein (2015) *Journal of Genetic Engineering and Biotechnology*, 13 (2), 145–155.
- 2.Borowitzka, M.A. and Borowitzka, L.J. (1988) *Micro-algal biotechnology*. Herausgegeben. Cambridge University Press, Cambridge, New York, New Rochella.
- 3.Avigad Vonshak and Amos Richmond (1988) *Mass production of the Blue-green alga Spirulina: An Overview, Biomass 15*, 233-247.
- 4.Kirensky LV, Terskov IA, Gitelson II, Lisovsky GM, Kovrov BG and Okladnikov YN.(1968) *Experimental biological life support system. I. Continuous cultivation of algae as a link of a closed ecosystem*. Life Sci Space Res. 6, 32-36.
- 5.A. Vonshak, S. Boussiba, A. Abeliovich and A. Richmond (1983) *Production of Spirulina Biomass: Maintenance of Monoalgal Culture Outdoors*, Biotechnology and Bioengineering, Vol. XXV, 341-349.
- 6.Vadiveloo A, Moheimani N.R, Alghamedi R, Cosgrove J.J, Alameh K and Parlevliet D. (2016) Sustainable cultivation of microalgae by an insulated glazed glass plate photobioreactor. *Biotechnol J*. 11(3), 363-74.
- 7.Habib MAB, Parvin, M, Huntington, T C and Hasan MR (2008) *A review on culture, production and use of Spirulina as food for humans and feeds for domestic animals and fish*, FAO Fisheries and Aquaculture Circular No. 1034, Rome, FAO, p.33.
- 8.Mori K, Ohya H, Matsumoto K, Furuune H, Isozaki K and Siekmeier P (1989) *Design for a bioreactor with sunlight supply and operations systems for use in the space environment*. Adv Space Res. 9 (8):161-168.